

Analysis of Genome Grimelysin-Containing Locus in the Genome of *Serratia grimesii* A2

Gilyazeva A., Toymentseva A., Mardanov A.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Springer Science+Business Media New York. *Serratia grimesii* strain A2 was isolated from buffered solution of actin by workers of Institute of Marine Biology, Vladivostok city [1]. This bacterium attracts special interest because of its ability to produce intracellular proteinase grimelysin which hydrolyzes actin in the single site and thus abolishes actin polymerization [1]. In addition to this well-characterized function, grimelysin is also required for invasion of bacteria into eukaryotic cells [1]. Here, we present the analysis of genome sequence of *S. grimesii* strain A2. The 5133.068 bp draft genome consists of 120 contigs with 4999 coding sequences, including 82 RNA genes. Genome sequencing and annotation revealed the presence of grimelysin gene in the operon with the gene of hypothetical protein of unknown function and the presence of regulatory gene upstream of it. In silico analysis of promoter sequences allowed the identification of putative binding sites for regulatory proteins. We anticipate that the *S. grimesii* strain A2 genome analysis will advance the current knowledge about grimelysin functions and regulation of its expression.

<http://dx.doi.org/10.1007/s12668-016-0330-x>

Keywords

Genome sequencing, Grimelysin, Locus, Operon, *Serratia grimesii*

References

- [1] Bozhokina, E. S., Tsaplina, O. A., Efremova, T. N., et al. (2011). Bacterial invasion of eukaryotic cells can be mediated by actin-hydrolysing metalloproteases grimelysin and protealysin. *Cell Biology International*, 35, 111–118. doi:10.1042/CBI20100314.
- [2] Mahlen, S. D. (2011). *Serratia* infections: from military experiments to current practice. *Clinical Microbiology Reviews*, 24, 755–791. doi:10.1128/CMR.00017-11.
- [3] Breidenstein, E. B., Janot, L., Strehmel, J., et al. (2012). The Lon protease is essential for full virulence in *Pseudomonas aeruginosa*. *PloS One*. doi:10.1371/journal.pone.0049123.
- [4] Stock, I., Grueger, T., Wiedemann, B. (2003). Natural antibiotic susceptibility of strains of *Serratia marcescens* and the *S. liquefaciens* complex: *S. liquefaciens sensu stricto*, *S. proteamaculans*, and *S. grimesii*. *International Journal of Antimicrobial Agents*, 22, 35–47.
- [5] Mardanov, A. M., Toymentseva, A. A., Gilyazeva, A. G. (2014). Draft genome sequence of *Serratia grimesii* strain A2. *Genome Announcements*. doi:10.1128/genomeA.00937-14.
- [6] Mosel, M., Li, L., Drlica, K., Zhao, X. (2013). Superoxide-mediated protection of *Escherichia coli* from antimicrobials. *Antimicrobial Agents and Chemotherapy*, 11, 5755–5759. doi:10.1128/AAC.00754-13.

- [7] Remes, B., Berghoff, B. A., Forstner, K. U., Klug, G. (2014). Role of oxygen and the OxyR protein in the response to iron limitation in *Rhodobacter sphaeroides*. *BMC Genomics*, 15, 794. doi:10.1186/1471-2164--5-794.
- [8] Nicoloff, H., Arsene-Ploetze, F., Malandain, C., et al. (2004). Two arginine repressors regulate arginine biosynthesis in *Lactobacillus plantarum*. *Journal of Bacteriology*, 186, 6059–6069.
- [9] Wei, Y., & Terns, M. P. (2016). CRISPR outsourcing: commissioning IHF for site-specific integration of foreign DNA at the CRISPR array. *Molecular Cell*, 62, 803–804. doi:10.1016/j.molcel.2016.06.004.
- [10] Solaiman, D. K., & Swingle, B. M. (2010). Isolation of novel *Pseudomonas syringae* promoters and functional characterization in polyhydroxyalkanoate-producing pseudomonads. *New Biotechnology*, 27, 1–9. doi:10.1016/j.nbt.2009.12.003.